

# On the curious commutativity of AMPD matrices

Adhemar Bultheel

Dept. Computer Science, KU Leuven

Leipzig, 15 February 2018

<http://nalag.cs.kuleuven.be/papers/ade/LEIPZIG18>

# Definition

$$\mathbf{AMPD} = \mathbf{AM} + \mathbf{D}$$

- $A, D \in \mathbb{C}^{n \times n}$  diagonal matrices

- $M = M_\pi = \prod_{k \in \pi}^{\rightarrow} G_k = G_{\pi_1} G_{\pi_2} \cdots G_{\pi_n},$

$\pi = (\pi_1, \pi_2, \dots, \pi_n)$  a permutation of  $(1, 2, \dots, n)$

- $G_k = \begin{bmatrix} I_{k-1} & & & \\ & \alpha_k & \beta_k & \\ & \gamma_k & \delta_k & \\ & & & I_{n-k-1} \end{bmatrix}, \quad k = 1, \dots, n-1$   
 $G_n = \begin{bmatrix} I_{n-1} & \\ & \alpha_n \end{bmatrix}.$

- Note that  $G_i$  and  $G_j$  if  $|i - j| \geq 2$ .
- But  $G_k G_{k+1} \neq G_{k+1} G_k$  in general.

$$\mathbf{AMPD} = \mathbf{AM} + \mathbf{D}$$

- $A, D \in \mathbb{C}^{n \times n}$  diagonal matrices

- $M = M_\pi = \prod_{k \in \pi}^{\rightarrow} G_k = G_{\pi_1} G_{\pi_2} \cdots G_{\pi_n},$

$\pi = (\pi_1, \pi_2, \dots, \pi_n)$  a permutation of  $(1, 2, \dots, n)$

- $G_k = \begin{bmatrix} I_{k-1} & & & \\ & \alpha_k & \beta_k & \\ & \gamma_k & \delta_k & \\ & & & I_{n-k-1} \end{bmatrix}, \quad k = 1, \dots, n-1$   
 $G_n = \begin{bmatrix} I_{n-1} & \\ & \alpha_n \end{bmatrix}.$

- Note that  $G_i$  and  $G_j$  if  $|i - j| \geq 2$ .
- But  $G_k G_{k+1} \neq G_{k+1} G_k$  in general.

# Definition

$$\mathbf{AMPD} = \mathbf{AM} + \mathbf{D}$$

- $A, D \in \mathbb{C}^{n \times n}$  diagonal matrices

- $M = M_\pi = \prod_{k \in \pi}^{\rightarrow} G_k = G_{\pi_1} G_{\pi_2} \cdots G_{\pi_n},$

$\pi = (\pi_1, \pi_2, \dots, \pi_n)$  a permutation of  $(1, 2, \dots, n)$

- $G_k = \begin{bmatrix} I_{k-1} & & & \\ & \alpha_k & \beta_k & \\ & \gamma_k & \delta_k & \\ & & & I_{n-k-1} \end{bmatrix}, \quad k = 1, \dots, n-1$   
 $G_n = \begin{bmatrix} I_{n-1} & \\ & \alpha_n \end{bmatrix}.$

- Note that  $G_i$  and  $G_j$  if  $|i - j| \geq 2$ .
- But  $G_k G_{k+1} \neq G_{k+1} G_k$  in general.

$$\mathbf{AMPD} = \mathbf{AM} + \mathbf{D}$$

- $A, D \in \mathbb{C}^{n \times n}$  diagonal matrices

- $M = M_\pi = \prod_{k \in \pi}^{\rightarrow} G_k = G_{\pi_1} G_{\pi_2} \cdots G_{\pi_n},$

$\pi = (\pi_1, \pi_2, \dots, \pi_n)$  a permutation of  $(1, 2, \dots, n)$

- $G_k = \begin{bmatrix} I_{k-1} & & & \\ & \alpha_k & \beta_k & \\ & \gamma_k & \delta_k & \\ & & & I_{n-k-1} \end{bmatrix}, \quad k = 1, \dots, n-1$

$$G_n = \begin{bmatrix} I_{n-1} & \\ & \alpha_n \end{bmatrix}.$$

- Note that  $G_i$  and  $G_j$  if  $|i - j| \geq 2$ .
- But  $G_k G_{k+1} \neq G_{k+1} G_k$  in general.

$$\mathbf{AMPD} = \mathbf{AM} + \mathbf{D}$$

- $A, D \in \mathbb{C}^{n \times n}$  diagonal matrices

- $M = M_\pi = \prod_{k \in \pi}^{\rightarrow} G_k = G_{\pi_1} G_{\pi_2} \cdots G_{\pi_n},$

$\pi = (\pi_1, \pi_2, \dots, \pi_n)$  a permutation of  $(1, 2, \dots, n)$

- $G_k = \begin{bmatrix} I_{k-1} & & & \\ & \alpha_k & \beta_k & \\ & \gamma_k & \delta_k & \\ & & & I_{n-k-1} \end{bmatrix}, \quad k = 1, \dots, n-1$

$$G_n = \begin{bmatrix} I_{n-1} & \\ & \alpha_n \end{bmatrix}.$$

- Note that  $G_i$  and  $G_j$  if  $|i - j| \geq 2$ .
- But  $G_k G_{k+1} \neq G_{k+1} G_k$  in general.

# Table of contents

- Definition AMPD and observation
- The proof
- Unitary case
- The origin OPUC
- Generalization (ORFUC)
- Rational AMPD = RAMPD

## Example 2

- $G_1 = \begin{bmatrix} \alpha_1 & \beta_1 & 0 \\ \gamma_1 & \delta_1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, G_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \alpha_2 & \beta_2 \\ 0 & \gamma_2 & \delta_2 \end{bmatrix},$

- $G_1 G_2 = \begin{bmatrix} \alpha_1 & \beta_1 \alpha_2 & \beta_1 \beta_2 \\ \gamma_1 & \delta_1 \alpha_2 & \delta_1 \beta_2 \\ 0 & \gamma_2 & \delta_2 \end{bmatrix} \neq G_2 G_1 = \begin{bmatrix} \alpha_1 & \beta_1 & 0 \\ \alpha_2 \gamma_1 & \alpha_2 \delta_1 & \beta_2 \\ \gamma_2 \gamma_1 & \gamma_2 \delta_1 & \delta_2 \end{bmatrix}$

BUT

- $\det(G_1 G_2) = \det(G_2 G_1)$
- Thus also  $\det(AG_1 G_2 + D) = \det(AG_2 G_1 + D)$
- $D \rightarrow D - \lambda I \Rightarrow \sigma(AG_1 G_1 + D) = \sigma(AG_2 G_1 + D)$



## Example 2

- $G_1 = \begin{bmatrix} \alpha_1 & \beta_1 & 0 \\ \gamma_1 & \delta_1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, G_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \alpha_2 & \beta_2 \\ 0 & \gamma_2 & \delta_2 \end{bmatrix},$

- $G_1 G_2 = \begin{bmatrix} \alpha_1 & \beta_1 \alpha_2 & \beta_1 \beta_2 \\ \gamma_1 & \delta_1 \alpha_2 & \delta_1 \beta_2 \\ 0 & \gamma_2 & \delta_2 \end{bmatrix} \neq G_2 G_1 = \begin{bmatrix} \alpha_1 & \beta_1 & 0 \\ \alpha_2 \gamma_1 & \alpha_2 \delta_1 & \beta_2 \\ \gamma_2 \gamma_1 & \gamma_2 \delta_1 & \delta_2 \end{bmatrix}$

BUT


- $\det(G_1 G_2) = \det(G_2 G_1)$
- Thus also  $\det(AG_1 G_2 + D) = \det(AG_2 G_1 + D)$
- $D \rightarrow D - \lambda I \Rightarrow \sigma(AG_1 G_1 + D) = \sigma(AG_2 G_1 + D)$

## Example 2

- $G_1 = \begin{bmatrix} \alpha_1 & \beta_1 & 0 \\ \gamma_1 & \delta_1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, G_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \alpha_2 & \beta_2 \\ 0 & \gamma_2 & \delta_2 \end{bmatrix},$

- $G_1 G_2 = \begin{bmatrix} \alpha_1 & \beta_1 \alpha_2 & \beta_1 \beta_2 \\ \gamma_1 & \delta_1 \alpha_2 & \delta_1 \beta_2 \\ 0 & \gamma_2 & \delta_2 \end{bmatrix} \neq G_2 G_1 = \begin{bmatrix} \alpha_1 & \beta_1 & 0 \\ \alpha_2 \gamma_1 & \alpha_2 \delta_1 & \beta_2 \\ \gamma_2 \gamma_1 & \gamma_2 \delta_1 & \delta_2 \end{bmatrix}$

**BUT**




- $\det(G_1 G_2) = \det(G_2 G_1)$
- Thus also  $\det(AG_1 G_2 + D) = \det(AG_2 G_1 + D)$
- $D \rightarrow D - \lambda I \Rightarrow \sigma(AG_1 G_1 + D) = \sigma(AG_2 G_1 + D)$

## Example 2

- $G_1 = \begin{bmatrix} \alpha_1 & \beta_1 & 0 \\ \gamma_1 & \delta_1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, G_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \alpha_2 & \beta_2 \\ 0 & \gamma_2 & \delta_2 \end{bmatrix},$

- $G_1 G_2 = \begin{bmatrix} \alpha_1 & \beta_1 \alpha_2 & \beta_1 \beta_2 \\ \gamma_1 & \delta_1 \alpha_2 & \delta_1 \beta_2 \\ 0 & \gamma_2 & \delta_2 \end{bmatrix} \neq G_2 G_1 = \begin{bmatrix} \alpha_1 & \beta_1 & 0 \\ \alpha_2 \gamma_1 & \alpha_2 \delta_1 & \beta_2 \\ \gamma_2 \gamma_1 & \gamma_2 \delta_1 & \delta_2 \end{bmatrix}$

**BUT**




- $\det(G_1 G_2) = \det(G_2 G_1)$
- Thus also  $\det(AG_1 G_2 + D) = \det(AG_2 G_1 + D)$
- $D \rightarrow D - \lambda I \Rightarrow \sigma(AG_1 G_1 + D) = \sigma(AG_2 G_1 + D)$

## Example 2

- $G_1 = \begin{bmatrix} \alpha_1 & \beta_1 & 0 \\ \gamma_1 & \delta_1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, G_2 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \alpha_2 & \beta_2 \\ 0 & \gamma_2 & \delta_2 \end{bmatrix},$

- $G_1 G_2 = \begin{bmatrix} \alpha_1 & \beta_1 \alpha_2 & \beta_1 \beta_2 \\ \gamma_1 & \delta_1 \alpha_2 & \delta_1 \beta_2 \\ 0 & \gamma_2 & \delta_2 \end{bmatrix} \neq G_2 G_1 = \begin{bmatrix} \alpha_1 & \beta_1 & 0 \\ \alpha_2 \gamma_1 & \alpha_2 \delta_1 & \beta_2 \\ \gamma_2 \gamma_1 & \gamma_2 \delta_1 & \delta_2 \end{bmatrix}$

**BUT**



- $\det(G_1 G_2) = \det(G_2 G_1)$
- Thus also  $\det(AG_1 G_2 + D) = \det(AG_2 G_1 + D)$
- $D \rightarrow D - \lambda I \Rightarrow \sigma(AG_1 G_1 + D) = \sigma(AG_2 G_1 + D)$

## Question


- Is in general  $\sigma(AM_\pi + D)$  independent of  $\pi$ ?

- Let's do some experiments
- ... and the result is...

- 
- Now prove it!!!

## Question

- Is in general  $\sigma(AM_\pi + D)$  independent of  $\pi$ ?




- Let's do some experiments
- ... and the result is...

- 
- Now prove it!!!

## Question

- Is in general  $\sigma(AM_\pi + D)$  independent of  $\pi$ ?




- Let's do some experiments
- ... and the result is...



- Now prove it!!!

## Question

- Is in general  $\sigma(AM_\pi + D)$  independent of  $\pi$ ?



- Let's do some experiments
- ... and the result is...


**YES!**

- 
- Now prove it!!!



## Question

- Is in general  $\sigma(AM_\pi + D)$  independent of  $\pi$ ?



- Let's do some experiments
- ... and the result is...

**YES!**

- 
- Now prove it!!!

# What does $M_\pi$ look like?

The matrix  $M$  is the product of a number of  $G$ -matrices

$$M_1 = \begin{array}{cccccc} & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & & & \\ & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & & & \\ & & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & & \\ & & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & & \\ & & & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & \\ & & & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & \\ & & & & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} \end{array} \quad \text{or} \quad M_2 = \begin{array}{cccccc} & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & & & \\ & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & & & \\ & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & & & \\ & & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & & \\ & & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & & \\ & & & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & \\ & & & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & \end{array} \quad \text{or} \quad M_3 = \begin{array}{cccccc} & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & & & \\ & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & & & \\ & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & & & \\ & & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & & \\ & & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & & \\ & & & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & \\ & & & & \begin{array}{c} \rightarrow \\ \leftarrow \end{array} & \end{array}$$



$M_1$  = upper Hessenberg matrix  $\pi = (1, 2, 3, 4, 5, 6)$

$M_1$  :  $\pi = (2, 1, 3, 4, 6, 5)$  or  $(2, 3, 1, 6, 4, 5)$  or ...

$M_3$  = CMV matrix  $\pi = (1, 3, 5, 2, 4, 6)$  or  $(5, 1, 3, 4, 2, 6)$ , or ...

# Shape of the matrix

general order vs. CMV



$\alpha$  upper Hessenberg  $\prod_k^{\rightarrow} G_k$

$\beta$  lower Hessenberg  $\prod_k^{\leftarrow} G_k$

CMV: alternate  $(G_1)(G_3 G_2)(G_5 G_4) \cdots = (G_1 G_3 \cdots)(G_2 G_4 \cdots)$



## Lemma

Let  $\mathbb{C}^{n \times n} \ni M' = \text{product of } (n-1) \text{ } G\text{-matrices.}$

$$M = \begin{bmatrix} M' & 0 \\ 0 & 1 \end{bmatrix} \text{ and } G = \begin{bmatrix} I_{n-1} & 0 & 0 \\ 0 & \alpha & \beta \\ 0 & \gamma & \delta \end{bmatrix}.$$

$A, D$  are  $(n+1) \times (n+1)$  diagonal matrices.

Then

$$\det(AMG + D) = \det(AGM + D)$$

and hence

$$\sigma(AMG + D) = \sigma(AGM + D).$$

If  $\det$  are the same then take  $D \rightarrow D - \lambda I_{n+1}$

This is the characteristic polynomial  $\Rightarrow$  the  $\sigma$  are the same.

## Lemma

Let  $\mathbb{C}^{n \times n} \ni M' = \text{product of } (n-1) \text{ } G\text{-matrices.}$

$$M = \begin{bmatrix} M' & 0 \\ 0 & 1 \end{bmatrix} \text{ and } G = \begin{bmatrix} I_{n-1} & 0 & 0 \\ 0 & \alpha & \beta \\ 0 & \gamma & \delta \end{bmatrix}.$$

$A, D$  are  $(n+1) \times (n+1)$  diagonal matrices.

Then

$$\det(AMG + D) = \det(AGM + D)$$

and hence

$$\sigma(AMG + D) = \sigma(AGM + D).$$

If  $\det$  are the same then take  $D \rightarrow D - \lambda I_{n+1}$

This is the characteristic polynomial  $\Rightarrow$  the  $\sigma$  are the same.

- Set  $M' = \left[ \begin{array}{c|c} M'' & \mathbf{c} \\ \hline \mathbf{r} & m \end{array} \right]$ ,  $A = \text{diag}(A'', a', a)$ ,  $D = \text{diag}(D'', d', d)$ .

$$AGM + D = \left[ \begin{array}{c|cc} A''M'' + D'' & A''\mathbf{c} & 0 \\ \hline a'\alpha\mathbf{r} & a'\alpha m + d' & a'\beta \\ a'\gamma\mathbf{r} & a\gamma & a\delta + d \end{array} \right]$$

$$AMG + D = \left[ \begin{array}{c|cc} A''M'' + D'' & \alpha A''\mathbf{c} & \beta A''\mathbf{c} \\ \hline a'\mathbf{r} & a'\alpha m + d' & a'\beta m \\ 0 & a\gamma & a\delta + d \end{array} \right]$$

- Set  $M' = \left[ \begin{array}{c|c} M'' & \mathbf{c} \\ \hline \mathbf{r} & m \end{array} \right]$ ,  $A = \text{diag}(A'', a', a)$ ,  $D = \text{diag}(D'', d', d)$ .

- 

$$AGM + D = \left[ \begin{array}{c|cc} A''M'' + D'' & A''\mathbf{c} & 0 \\ \hline a'\alpha\mathbf{r} & a'\alpha m + d' & a'\beta \\ a\gamma\mathbf{r} & a\gamma & a\delta + d \end{array} \right]$$

$$AMG + D = \left[ \begin{array}{c|cc} A''M'' + D'' & \alpha A''\mathbf{c} & \beta A''\mathbf{c} \\ \hline a'\mathbf{r} & a'\alpha m + d' & a'\beta m \\ 0 & a\gamma & a\delta + d \end{array} \right]$$



- work out ...
- regroup ...

- 
- $\det(AGM + D) = \det(AMG + D) =$   
 $[aa' \det G + a'd\alpha] \det \tilde{M} + (a\delta + d)d' \det(A''M'' + D'')$   
 $\tilde{M} = \text{diag}(A'', 1)M' + \text{diag}(D'', 0)$
- Thus determinants (hence the spectra) are the same.

- work out ...
- regroup ...

- 
- $\det(AGM + D) = \det(AMG + D) =$   
 $[aa' \det G + a'd\alpha] \det \tilde{M} + (a\delta + d)d' \det(A''M'' + D'')$   
 $\tilde{M} = \text{diag}(A'', 1)M' + \text{diag}(D'', 0)$
- Thus determinants (hence the spectra) are the same.

- work out ...
- regroup ...



- 
- $\det(AGM + D) = \det(AMG + D) =$   
 $[aa' \det G + a'd\alpha] \det \tilde{M} + (a\delta + d)d' \det(A''M'' + D'')$   
 $\tilde{M} = \text{diag}(A'', 1)M' + \text{diag}(D'', 0)$
- Thus determinants (hence the spectra) are the same.

- work out ...
- regroup ...



- 
- $\det(AGM + D) = \det(AMG + D) =$   
 $[aa' \det G + a'd\alpha] \det \tilde{M} + (a\delta + d)d' \det(A''M'' + D'')$   
 $\tilde{M} = \text{diag}(A'', 1)M' + \text{diag}(D'', 0)$
- Thus determinants (hence the spectra) are the same.

- work out ...
- regroup ...



- 
- $\det(AGM + D) = \det(AMG + D) =$   
 $[aa' \det G + a'd\alpha] \det \tilde{M} + (a\delta + d)d' \det(A''M'' + D'')$   
 $\tilde{M} = \text{diag}(A'', 1)M' + \text{diag}(D'', 0)$
- Thus determinants (hence the spectra) are the same.

Projection:  $P_n^N = [I_n \ 0] \in \mathbb{C}^{n \times N}$ , (e.g.  $P = P_n^{n+1} \Rightarrow M' = PMP^*$ )

## Theorem

$A, D$  diagonal of size  $n + 1$

$\pi = (\pi_1, \dots, \pi_n)$  = permutation of  $(1, \dots, n)$

$M_\pi = G_{\pi_1} G_{\pi_2} \cdots G_{\pi_n} \in \mathbb{C}^{(n+1) \times (n+1)}$ ,  $P = P_n^{n+1}$

Then  $\det(P(AM_\pi + D)P^*)$  independent of  $\pi$

hence also  $\sigma(P(AM_\pi + D)P^*)$  independent of  $\pi$

OK for  $n = 2$  (see Example 2)

# Proof $n = 2$

Recall Example 2:

$$G_1 G_2 = \begin{bmatrix} \alpha_1 & \beta_1 \alpha_2 & \beta_1 \beta_2 \\ \gamma_1 & \delta_1 \alpha_2 & \delta_1 \beta_2 \\ 0 & \gamma_2 & \delta_2 \end{bmatrix} \neq G_2 G_1 = \begin{bmatrix} \alpha_1 & \beta_1 & 0 \\ \alpha_2 \gamma_1 & \alpha_2 \delta_1 & \beta_2 \\ \gamma_2 \gamma_1 & \gamma_2 \delta_1 & \delta_2 \end{bmatrix}$$

$$AG_1 G_2 + D = \left[ \begin{array}{cc|c} a_1 \alpha_1 + d_1 & a_1 \beta_1 \alpha_2 & a_1 \beta_1 \beta_2 \\ a_2 \gamma_1 & a_2 \delta_1 \alpha_2 + d_2 & a_2 \delta_1 \beta_2 \\ \hline 0 & a_3 \gamma_2 & a_3 \delta_2 + d_3 \end{array} \right]$$

$$AG_2 G_1 + D = \left[ \begin{array}{cc|c} a_1 \alpha_1 + d_1 & a_1 \beta_1 & 0 \\ a_2 \alpha_2 \gamma_1 & a_2 \alpha_2 \delta_1 + d_2 & a_2 \beta_2 \\ \hline a_3 \gamma_2 \gamma_1 & a_3 \gamma_2 \delta_1 & a_3 \delta_2 + d_3 \end{array} \right]$$

$$\text{hence } \det(P(AG_1 G_2 + D)P^*) = \det(P(AG_2 G_1 + D)P^*)$$

# Proof Induction step

- Set  $\pi' = \text{permutation of } (1, \dots, n-1)$   
Then  $M_\pi = G_n M_{\pi'}$  or  $M_{\pi'} G_n$ .
- Suppose  $M_\pi = G_n M_{\pi'}$ :

$$P(M_\pi)P^* = P \left[ \begin{array}{c|c|c} M'' & \mathbf{c}_{n-1} & 0 \\ \hline \alpha_n \mathbf{r}_{n-1} & \alpha_n m_{n-1} & \beta_n \\ \hline \gamma_n \mathbf{r}_{n-1} & \gamma_n m_{n-1} & \delta_n \end{array} \right] P^* = P \left[ \begin{array}{c|c|c} M'' & \mathbf{c}_{n-1} & 0 \\ \hline \alpha_n \mathbf{r}_{n-1} & \alpha_n m_{n-1} & 0 \\ \hline 0 & 0 & 1 \end{array} \right] P^*$$

- $\det(P(AM_\pi + D)P^*) = \det[(PAM_\pi P^*) + PDP^*] = \det(\hat{A}\hat{M} + \hat{D})$   
where  $\hat{A} = \text{diag}(1, \dots, 1, \alpha_n)PAP^*$ ,  $\hat{D} = PDP^*$ ,  $\hat{M} = PM_{\pi'}P^*$
- $\hat{A}\hat{M} + \hat{D} \in \mathbb{C}^{n \times n}$  is an AMPD matrix with  $n-1$  G-factors.
- Similarly for  $M_\pi = M_{\pi'} G_n$ .



# Proof Induction step

- Set  $\pi' = \text{permutation of } (1, \dots, n-1)$   
Then  $M_\pi = G_n M_{\pi'}$  or  $M_{\pi'} G_n$ .
- Suppose  $M_\pi = G_n M_{\pi'}$ :

$$P(M_\pi)P^* = P \left[ \begin{array}{c|c|c} M'' & \mathbf{c}_{n-1} & 0 \\ \hline \alpha_n \mathbf{r}_{n-1} & \alpha_n \mathbf{m}_{n-1} & \beta_n \\ \hline \gamma_n \mathbf{r}_{n-1} & \gamma_n \mathbf{m}_{n-1} & \delta_n \end{array} \right] P^* = P \left[ \begin{array}{c|c|c} M'' & \mathbf{c}_{n-1} & 0 \\ \hline \alpha_n \mathbf{r}_{n-1} & \alpha_n \mathbf{m}_{n-1} & 0 \\ \hline 0 & 0 & 1 \end{array} \right] P^*$$

- $\det(P(AM_\pi + D)P^*) = \det[(PAM_\pi P^*) + PDP^*] = \det(\hat{A}\hat{M} + \hat{D})$   
where  $\hat{A} = \text{diag}(1, \dots, 1, \alpha_n)PAP^*$ ,  $\hat{D} = PDP^*$ ,  $\hat{M} = PM_{\pi'}P^*$
- $\hat{A}\hat{M} + \hat{D} \in \mathbb{C}^{n \times n}$  is an AMPD matrix with  $n-1$  G-factors.
- Similarly for  $M_\pi = M_{\pi'} G_n$ .

# Proof Induction step

- Set  $\pi' = \text{permutation of } (1, \dots, n-1)$   
Then  $M_\pi = G_n M_{\pi'}$  or  $M_{\pi'} G_n$ .
- Suppose  $M_\pi = G_n M_{\pi'}$ :

$$P(M_\pi)P^* = P \left[ \begin{array}{c|c|c} M'' & \mathbf{c}_{n-1} & 0 \\ \hline \alpha_n \mathbf{r}_{n-1} & \alpha_n m_{n-1} & \beta_n \\ \hline \gamma_n \mathbf{r}_{n-1} & \gamma_n m_{n-1} & \delta_n \end{array} \right] P^* = P \left[ \begin{array}{c|c|c} M'' & \mathbf{c}_{n-1} & 0 \\ \hline \alpha_n \mathbf{r}_{n-1} & \alpha_n m_{n-1} & 0 \\ \hline 0 & 0 & 1 \end{array} \right] P^*$$

- $\det(P(AM_\pi + D)P^*) = \det[(PAM_\pi P^*) + PDP^*] = \det(\hat{A}\hat{M} + \hat{D})$   
where  $\hat{A} = \text{diag}(1, \dots, 1, \alpha_n)PAP^*$ ,  $\hat{D} = PDP^*$ ,  $\hat{M} = PM_{\pi'}P^*$
- $\hat{A}\hat{M} + \hat{D} \in \mathbb{C}^{n \times n}$  is an AMPD matrix with  $n-1$  G-factors.
- Similarly for  $M_\pi = M_{\pi'} G_n$ .

# Proof Induction step

- Set  $\pi' = \text{permutation of } (1, \dots, n-1)$   
Then  $M_\pi = G_n M_{\pi'}$  or  $M_{\pi'} G_n$ .
- Suppose  $M_\pi = G_n M_{\pi'}$ :

$$P(M_\pi)P^* = P \left[ \begin{array}{c|c|c} M'' & \mathbf{c}_{n-1} & 0 \\ \hline \alpha_n \mathbf{r}_{n-1} & \alpha_n \mathbf{m}_{n-1} & \beta_n \\ \hline \gamma_n \mathbf{r}_{n-1} & \gamma_n \mathbf{m}_{n-1} & \delta_n \end{array} \right] P^* = P \left[ \begin{array}{c|c|c} M'' & \mathbf{c}_{n-1} & 0 \\ \hline \alpha_n \mathbf{r}_{n-1} & \alpha_n \mathbf{m}_{n-1} & 0 \\ \hline 0 & 0 & 1 \end{array} \right] P^*$$

- $\det(P(AM_\pi + D)P^*) = \det[(PAM_\pi P^*) + PDP^*] = \det(\hat{A}\hat{M} + \hat{D})$   
where  $\hat{A} = \text{diag}(1, \dots, 1, \alpha_n)PAP^*$ ,  $\hat{D} = PDP^*$ ,  $\hat{M} = PM_{\pi'}P^*$
- $\hat{A}\hat{M} + \hat{D} \in \mathbb{C}^{n \times n}$  is an AMPD matrix with  $n-1$  G-factors.
- Similarly for  $M_\pi = M_{\pi'} G_n$ .

# Proof Induction step

- Set  $\pi' = \text{permutation of } (1, \dots, n-1)$   
Then  $M_\pi = G_n M_{\pi'}$  or  $M_{\pi'} G_n$ .
- Suppose  $M_\pi = G_n M_{\pi'}$ :

$$P(M_\pi)P^* = P \left[ \begin{array}{c|c|c} M'' & \mathbf{c}_{n-1} & 0 \\ \hline \alpha_n \mathbf{r}_{n-1} & \alpha_n m_{n-1} & \beta_n \\ \hline \gamma_n \mathbf{r}_{n-1} & \gamma_n m_{n-1} & \delta_n \end{array} \right] P^* = P \left[ \begin{array}{c|c|c} M'' & \mathbf{c}_{n-1} & 0 \\ \hline \alpha_n \mathbf{r}_{n-1} & \alpha_n m_{n-1} & 0 \\ \hline 0 & 0 & 1 \end{array} \right] P^*$$

- $\det(P(AM_\pi + D)P^*) = \det[(PAM_\pi P^*) + PDP^*] = \det(\hat{A}\hat{M} + \hat{D})$   
where  $\hat{A} = \text{diag}(1, \dots, 1, \alpha_n)PAP^*$ ,  $\hat{D} = PDP^*$ ,  $\hat{M} = PM_{\pi'}P^*$
- $\hat{A}\hat{M} + \hat{D} \in \mathbb{C}^{n \times n}$  is an AMPD matrix with  $n-1$  G-factors.
- Similarly for  $M_\pi = M_{\pi'} G_n$ .

## Corollary

$A, D$  diagonal of size  $n + 1$

$\pi = (\pi_1, \dots, \pi_n) = \text{permutation of } (1, \dots, n)$

$M_\pi = G_{\pi_1} G_{\pi_2} \cdots G_{\pi_n} \in \mathbb{C}^{(n+1) \times (n+1)}$

Then  $\det(AM_\pi + D)$  independent of  $\pi$

hence also  $\sigma(AM_\pi + D)$  independent of  $\pi$

Proof: Use  $(n + 1)$   $G$ -factors with  $G_{n+1} = I_{n+2}$ .

Apply the previous theorem:

the projection  $P$  leaves the product of  $n$   $G$ -factors.

## Corollary

$A, D$  diagonal of size  $n + 1$

$\pi = (\pi_1, \dots, \pi_n)$  = permutation of  $(1, \dots, n)$

$M_\pi = G_{\pi_1} G_{\pi_2} \cdots G_{\pi_n} \in \mathbb{C}^{(n+1) \times (n+1)}$

Then  $\det(AM_\pi + D)$  independent of  $\pi$

hence also  $\sigma(AM_\pi + D)$  independent of  $\pi$

Proof: Use  $(n + 1)$   $G$ -factors with  $G_{n+1} = I_{n+2}$ .

Apply the previous theorem:

the projection  $P$  leaves the product of  $n$   $G$ -factors.



# Unitary case

- The eigenvalues are independent of  $\pi$  but the eigenvectors change
- However, suppose all  $G_k$  are unitary and do some more experiments.

- ...

- $M_\pi = V_\pi \Lambda V_\pi^*$

The eigenvalues (in  $\Lambda$ ) do not depend on  $\pi$

The eigenvalues are all on  $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$

The matrix of eigenvectors is unitary  $V_\pi^* V_\pi = I_{n+1}$

The absolute values  $|V_\pi| = [|v_{ij}^\pi|]_{i,j=1}^{n+1}$  do not depend on  $\pi$

- To prove this  $\rightarrow$  go to the origin of the problem:  
Orthogonal Polynomials (OPUC)

# Unitary case

- The eigenvalues are independent of  $\pi$  but the eigenvectors change
- However, suppose all  $G_k$  are unitary and do some more experiments.

• ...

- $M_\pi = V_\pi \Lambda V_\pi^*$

The eigenvalues (in  $\Lambda$ ) do not depend on  $\pi$

The eigenvalues are all on  $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$

The matrix of eigenvectors is unitary  $V_\pi^* V_\pi = I_{n+1}$


The absolute values  $|V_\pi| = [|v_{ij}^\pi|]_{i,j=1}^{n+1}$  do not depend on  $\pi$

- To prove this  $\rightarrow$  go to the origin of the problem:  
Orthogonal Polynomials (OPUC)



# Unitary case

- The eigenvalues are independent of  $\pi$  but the eigenvectors change
- However, suppose all  $G_k$  are unitary and do some more experiments.



• ...

- $M_\pi = V_\pi \Lambda V_\pi^*$

The eigenvalues (in  $\Lambda$ ) do not depend on  $\pi$

The eigenvalues are all on  $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$

The matrix of eigenvectors is unitary  $V_\pi^* V_\pi = I_{n+1}$


The absolute values  $|V_\pi| = [|v_{ij}^\pi|]_{i,j=1}^{n+1}$  do not depend on  $\pi$

- To prove this  $\rightarrow$  go to the origin of the problem:

Orthogonal Polynomials (OPUC)

# Unitary case

- The eigenvalues are independent of  $\pi$  but the eigenvectors change
- However, suppose all  $G_k$  are unitary and do some more experiments.



- ...

- $M_\pi = V_\pi \Lambda V_\pi^*$

The eigenvalues (in  $\Lambda$ ) do not depend on  $\pi$

The eigenvalues are all on  $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$

The matrix of eigenvectors is unitary  $V_\pi^* V_\pi = I_{n+1}$


The absolute values  $|V_\pi| = [|v_{ij}^\pi|]_{i,j=1}^{n+1}$  do not depend on  $\pi$

- To prove this  $\rightarrow$  go to the origin of the problem:

Orthogonal Polynomials (OPUC)

# Unitary case

- The eigenvalues are independent of  $\pi$  but the eigenvectors change
- However, suppose all  $G_k$  are unitary and do some more experiments.



- ...

- $M_\pi = V_\pi \Lambda V_\pi^*$

The eigenvalues (in  $\Lambda$ ) do not depend on  $\pi$

The eigenvalues are all on  $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$

The matrix of eigenvectors is unitary  $V_\pi^* V_\pi = I_{n+1}$

The absolute values  $|V_\pi| = [|v_{ij}^\pi|]_{i,j=1}^{n+1}$  do not depend on  $\pi$

- To prove this  $\rightarrow$  go to the origin of the problem:

**Orthogonal Polynomials** (OPUC)

# Szegő polynomials

- $\langle f, g \rangle = \int_{\mathbb{T}} \overline{f(z)} g(z) \mu(dz), \quad \mu(\mathbb{T}) = 1$
- $[1, z, z^2, \dots] \rightarrow$  orthonormalize  $\rightarrow \Phi = [\phi_0, \phi_1, \dots]$  then

$$z\Phi(z) = \Phi(z)\mathcal{G}, \quad \mathcal{G} = G_0 G_1 G_2 \cdots, \quad G_k = \begin{bmatrix} I_k & & & \\ & -\delta_k & \eta_k & \\ & \eta_k & \bar{\delta}_k & \\ & & & I_\infty \end{bmatrix}$$

$\delta_k \in \mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ ,  $\eta_k = \sqrt{1 - |\delta_k|^2} \Rightarrow G_k$  unitary  
 $\mathcal{G}$  upper Hessenberg

- Truncate:  $z\Phi_n(z) = \Phi_n(z)\mathcal{G}_n + [0, \dots, 0, c\phi_{n+1}]$   
with  $P = P_{n+1}^\infty$ :  $\Phi_n = \Phi P^*$  and  $\mathcal{G}_n = P\mathcal{G}P^*$ .
- If  $z = z_i$  is zero of  $\phi_{n+1}$ ,  
then  $z_i \in \sigma(\mathcal{G}_n)$  and  
 $\Phi_n(z_i)$  is a corresponding (left) eigenvector.

# Szegő polynomials

- $\langle f, g \rangle = \int_{\mathbb{T}} \overline{f(z)} g(z) \mu(dz), \quad \mu(\mathbb{T}) = 1$
- $[1, z, z^2, \dots] \rightarrow$  orthonormalize  $\rightarrow \Phi = [\phi_0, \phi_1, \dots]$  then

$$z\Phi(z) = \Phi(z)\mathcal{G}, \quad \mathcal{G} = G_0 G_1 G_2 \cdots, \quad G_k = \begin{bmatrix} I_k & & & \\ & -\delta_k & \eta_k & \\ & \eta_k & \bar{\delta}_k & \\ & & & I_\infty \end{bmatrix}$$

$\delta_k \in \mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ ,  $\eta_k = \sqrt{1 - |\delta_k|^2} \Rightarrow G_k$  unitary  
 $\mathcal{G}$  upper Hessenberg

- Truncate:  $z\Phi_n(z) = \Phi_n(z)\mathcal{G}_n + [0, \dots, 0, c\phi_{n+1}]$   
with  $P = P_{n+1}^\infty$ :  $\Phi_n = \Phi P^*$  and  $\mathcal{G}_n = P\mathcal{G}P^*$ .
- If  $z = z_i$  is zero of  $\phi_{n+1}$ ,  
then  $z_i \in \sigma(\mathcal{G}_n)$  and  
 $\Phi_n(z_i)$  is a corresponding (left) eigenvector.

# Szegő polynomials

- $\langle f, g \rangle = \int_{\mathbb{T}} \overline{f(z)} g(z) \mu(dz), \quad \mu(\mathbb{T}) = 1$
- $[1, z, z^2, \dots] \rightarrow$  orthonormalize  $\rightarrow \Phi = [\phi_0, \phi_1, \dots]$  then

$$z\Phi(z) = \Phi(z)\mathcal{G}, \quad \mathcal{G} = G_0 G_1 G_2 \cdots, \quad G_k = \begin{bmatrix} I_k & & & \\ & -\delta_k & \eta_k & \\ & \eta_k & \bar{\delta}_k & \\ & & & I_\infty \end{bmatrix}$$

$\delta_k \in \mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ ,  $\eta_k = \sqrt{1 - |\delta_k|^2} \Rightarrow G_k$  unitary  
 $\mathcal{G}$  upper Hessenberg

- Truncate:  $z\Phi_n(z) = \Phi_n(z)\mathcal{G}_n + [0, \dots, 0, c\phi_{n+1}]$   
with  $P = P_{n+1}^\infty$ :  $\Phi_n = \Phi P^*$  and  $\mathcal{G}_n = P\mathcal{G}P^*$ .
- If  $z = z_i$  is zero of  $\phi_{n+1}$ ,  
then  $z_i \in \sigma(\mathcal{G}_n)$  and  
 $\Phi_n(z_i)$  is a corresponding (left) eigenvector.

# Szegő polynomials

- $\langle f, g \rangle = \int_{\mathbb{T}} \overline{f(z)} g(z) \mu(dz), \quad \mu(\mathbb{T}) = 1$
- $[1, z, z^2, \dots] \rightarrow$  orthonormalize  $\rightarrow \Phi = [\phi_0, \phi_1, \dots]$  then

$$z\Phi(z) = \Phi(z)\mathcal{G}, \quad \mathcal{G} = G_0 G_1 G_2 \cdots, \quad G_k = \begin{bmatrix} I_k & & & \\ & -\delta_k & \eta_k & \\ & \eta_k & \bar{\delta}_k & \\ & & & I_\infty \end{bmatrix}$$

$\delta_k \in \mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$ ,  $\eta_k = \sqrt{1 - |\delta_k|^2} \Rightarrow G_k$  unitary  
 $\mathcal{G}$  upper Hessenberg

- Truncate:  $z\Phi_n(z) = \Phi_n(z)\mathcal{G}_n + [0, \dots, 0, c\phi_{n+1}]$   
with  $P = P_{n+1}^\infty$ :  $\Phi_n = \Phi P^*$  and  $\mathcal{G}_n = P\mathcal{G}P^*$ .
- If  $z = z_i$  is zero of  $\phi_{n+1}$ ,  
then  $z_i \in \sigma(\mathcal{G}_n)$  and  
 $\Phi_n(z_i)$  is a corresponding (left) eigenvector.

- $\sigma(\mathcal{G}_n)$  independent of the order in which the  $\{G_k\}_{k=0}^{n-1}$  are multiplied
- CMV reduces the upper Hessenberg to a CMV (5-diagonal) matrix  
 $\mathcal{G} = G_0 G_1 G_2 G_3 G_4 G_5 \cdots \rightarrow \tilde{\mathcal{G}} = (G_0 G_2 G_4 \cdots)(G_1 G_3 G_5 \cdots)$
- **Unitary Truncation**  
If  $\mathcal{G}$  is unitary,  $\nrightarrow \mathcal{G}_n$  unitary
- Problem is that  $G_n = \begin{bmatrix} -\delta_n & \eta_n \\ \eta_n & \bar{\delta}_n \end{bmatrix}$   
truncated to  $G_n = [-\delta_n]$ , and  $\delta_n \in \mathbb{D}$ .
- To make this unitary choose  $\delta_n \in \mathbb{T}$
- then  $\mathcal{G}_n$  is unitary  $\Rightarrow$  eigenvalues on  $\mathbb{T}$
- zeros of  $\phi_{n+1}$  (paraorthogonal) are on  $\mathbb{T}$  and simple



- $\sigma(\mathcal{G}_n)$  independent of the order in which the  $\{G_k\}_{k=0}^{n-1}$  are multiplied
- CMV reduces the upper Hessenberg to a CMV (5-diagonal) matrix  
 $\mathcal{G} = G_0 G_1 G_2 G_3 G_4 G_5 \cdots \rightarrow \tilde{\mathcal{G}} = (G_0 G_2 G_4 \cdots)(G_1 G_3 G_5 \cdots)$

- **Unitary Truncation**

If  $\mathcal{G}$  is unitary,  $\nrightarrow \mathcal{G}_n$  unitary

- Problem is that  $G_n = \begin{bmatrix} -\delta_n & \eta_n \\ \eta_n & \bar{\delta}_n \end{bmatrix}$   
truncated to  $G_n = [-\delta_n]$ , and  $\delta_n \in \mathbb{D}$ .
- To make this unitary choose  $\delta_n \in \mathbb{T}$
- then  $\mathcal{G}_n$  is unitary  $\Rightarrow$  eigenvalues on  $\mathbb{T}$
- zeros of  $\phi_{n+1}$  (paraorthogonal) are on  $\mathbb{T}$  and simple

- $\sigma(\mathcal{G}_n)$  independent of the order in which the  $\{G_k\}_{k=0}^{n-1}$  are multiplied
- CMV reduces the upper Hessenberg to a CMV (5-diagonal) matrix  
 $\mathcal{G} = G_0 G_1 G_2 G_3 G_4 G_5 \cdots \rightarrow \tilde{\mathcal{G}} = (G_0 G_2 G_4 \cdots)(G_1 G_3 G_5 \cdots)$

- **Unitary Truncation**

If  $\mathcal{G}$  is unitary,  $\nrightarrow \mathcal{G}_n$  unitary

- Problem is that  $G_n = \begin{bmatrix} -\delta_n & \eta_n \\ \eta_n & \bar{\delta}_n \end{bmatrix}$   
truncated to  $G_n = [-\delta_n]$ , and  $\delta_n \in \mathbb{D}$ .
- To make this unitary choose  $\delta_n \in \mathbb{T}$
- then  $\mathcal{G}_n$  is unitary  $\Rightarrow$  eigenvalues on  $\mathbb{T}$
- zeros of  $\phi_{n+1}$  (paraorthogonal) are on  $\mathbb{T}$  and simple

- $\sigma(\mathcal{G}_n)$  independent of the order in which the  $\{G_k\}_{k=0}^{n-1}$  are multiplied
- CMV reduces the upper Hessenberg to a CMV (5-diagonal) matrix  $\mathcal{G} = G_0 G_1 G_2 G_3 G_4 G_5 \cdots \rightarrow \tilde{\mathcal{G}} = (G_0 G_2 G_4 \cdots)(G_1 G_3 G_5 \cdots)$

- **Unitary Truncation**

If  $\mathcal{G}$  is unitary,  $\nrightarrow \mathcal{G}_n$  unitary

- Problem is that  $G_n = \begin{bmatrix} -\delta_n & \eta_n \\ \eta_n & \bar{\delta}_n \end{bmatrix}$   
truncated to  $G_n = [-\delta_n]$ , and  $\delta_n \in \mathbb{D}$ .
- To make this unitary choose  $\delta_n \in \mathbb{T}$
- then  $\mathcal{G}_n$  is unitary  $\Rightarrow$  eigenvalues on  $\mathbb{T}$
- zeros of  $\phi_{n+1}$  (paraorthogonal) are on  $\mathbb{T}$  and simple

- $\sigma(\mathcal{G}_n)$  independent of the order in which the  $\{G_k\}_{k=0}^{n-1}$  are multiplied
- CMV reduces the upper Hessenberg to a CMV (5-diagonal) matrix  $\mathcal{G} = G_0 G_1 G_2 G_3 G_4 G_5 \cdots \rightarrow \tilde{\mathcal{G}} = (G_0 G_2 G_4 \cdots)(G_1 G_3 G_5 \cdots)$

- **Unitary Truncation**

If  $\mathcal{G}$  is unitary,  $\nrightarrow \mathcal{G}_n$  unitary

- Problem is that  $G_n = \begin{bmatrix} -\delta_n & \eta_n \\ \eta_n & \bar{\delta}_n \end{bmatrix}$   
truncated to  $G_n = [-\delta_n]$ , and  $\delta_n \in \mathbb{D}$ .
- To make this unitary choose  $\delta_n \in \mathbb{T}$
- then  $\mathcal{G}_n$  is unitary  $\Rightarrow$  eigenvalues on  $\mathbb{T}$
- zeros of  $\phi_{n+1}$  (paraorthogonal) are on  $\mathbb{T}$  and simple

- $\sigma(\mathcal{G}_n)$  independent of the order in which the  $\{G_k\}_{k=0}^{n-1}$  are multiplied
- CMV reduces the upper Hessenberg to a CMV (5-diagonal) matrix  $\mathcal{G} = G_0 G_1 G_2 G_3 G_4 G_5 \cdots \rightarrow \tilde{\mathcal{G}} = (G_0 G_2 G_4 \cdots)(G_1 G_3 G_5 \cdots)$

- **Unitary Truncation**

If  $\mathcal{G}$  is unitary,  $\nrightarrow \mathcal{G}_n$  unitary

- Problem is that  $G_n = \begin{bmatrix} -\delta_n & \eta_n \\ \eta_n & \bar{\delta}_n \end{bmatrix}$   
truncated to  $G_n = [-\delta_n]$ , and  $\delta_n \in \mathbb{D}$ .
- To make this unitary choose  $\delta_n \in \mathbb{T}$
- then  $\mathcal{G}_n$  is unitary  $\Rightarrow$  eigenvalues on  $\mathbb{T}$
- zeros of  $\phi_{n+1}$  (paraorthogonal) are on  $\mathbb{T}$  and simple

- $\sigma(\mathcal{G}_n)$  independent of the order in which the  $\{G_k\}_{k=0}^{n-1}$  are multiplied
- CMV reduces the upper Hessenberg to a CMV (5-diagonal) matrix  
 $\mathcal{G} = G_0 G_1 G_2 G_3 G_4 G_5 \cdots \rightarrow \tilde{\mathcal{G}} = (G_0 G_2 G_4 \cdots)(G_1 G_3 G_5 \cdots)$

- **Unitary Truncation**

If  $\mathcal{G}$  is unitary,  $\nrightarrow \mathcal{G}_n$  unitary

- Problem is that  $G_n = \begin{bmatrix} -\delta_n & \eta_n \\ \eta_n & \bar{\delta}_n \end{bmatrix}$   
truncated to  $G_n = [-\delta_n]$ , and  $\delta_n \in \mathbb{D}$ .
- To make this unitary choose  $\delta_n \in \mathbb{T}$
- then  $\mathcal{G}_n$  is unitary  $\Rightarrow$  eigenvalues on  $\mathbb{T}$
- zeros of  $\phi_{n+1}$  (paraorthogonal) are on  $\mathbb{T}$  and simple

## Note

- In  $M_\pi = V_\pi^* \Lambda V_\pi$  we assumed  $V_\pi^* V_\pi = I_{n+1}$
- Here  $\mathcal{G}_n = \mathcal{V}_n^* \Lambda \mathcal{V}_n$  but here  $\mathcal{V}_n^* \mathcal{V}_n \neq I_{n+1}$

$$\mathcal{V}_n = \begin{bmatrix} \Phi_n(z_0) \\ \Phi_n(z_1) \\ \vdots \\ \Phi_n(z_n) \end{bmatrix} = \begin{bmatrix} \phi_0 & \phi_1(z_0) & \cdots & \phi_n(z_0) \\ \phi_0 & \phi_1(z_1) & \cdots & \phi_n(z_1) \\ \vdots & \vdots & & \vdots \\ \phi_0 & \phi_1(z_n) & \cdots & \phi_n(z_n) \end{bmatrix}$$

because  $\Phi_n(z_i) = [\phi_0, \phi_1(z_i), \dots, \phi_n(z_i)]$  and  $\phi_0 = 1$ .

- But by renormalization:

$$V_n = N_n \mathcal{V}_n, \quad N_n = \text{diag}(\|\Phi_n(z_i)\|^{-1} : i = 0, \dots, n)$$

then  $V_n^* V_n = I_{n+1}$ .

## Note

- In  $M_\pi = V_\pi^* \Lambda V_\pi$  we assumed  $V_\pi^* V_\pi = I_{n+1}$
- Here  $\mathcal{G}_n = \mathcal{V}_n^* \Lambda \mathcal{V}_n$  but here  $\mathcal{V}_n^* \mathcal{V}_n \neq I_{n+1}$

$$\mathcal{V}_n = \begin{bmatrix} \Phi_n(z_0) \\ \Phi_n(z_1) \\ \vdots \\ \Phi_n(z_n) \end{bmatrix} = \begin{bmatrix} \phi_0 & \phi_1(z_0) & \cdots & \phi_n(z_0) \\ \phi_0 & \phi_1(z_1) & \cdots & \phi_n(z_1) \\ \vdots & \vdots & & \vdots \\ \phi_0 & \phi_1(z_n) & \cdots & \phi_n(z_n) \end{bmatrix}$$

because  $\Phi_n(z_i) = [\phi_0, \phi_1(z_i), \dots, \phi_n(z_i)]$  and  $\phi_0 = 1$ .

- But by renormalization:

$$V_n = N_n \mathcal{V}_n, \quad N_n = \text{diag}(\|\Phi_n(z_i)\|^{-1} : i = 0, \dots, n)$$

then  $V_n^* V_n = I_{n+1}$ .



## Note

- In  $M_\pi = V_\pi^* \Lambda V_\pi$  we assumed  $V_\pi^* V_\pi = I_{n+1}$
- Here  $\mathcal{G}_n = \mathcal{V}_n^* \Lambda \mathcal{V}_n$  but here  $\mathcal{V}_n^* \mathcal{V}_n \neq I_{n+1}$

$$\mathcal{V}_n = \begin{bmatrix} \Phi_n(z_0) \\ \Phi_n(z_1) \\ \vdots \\ \Phi_n(z_n) \end{bmatrix} = \begin{bmatrix} \phi_0 & \phi_1(z_0) & \cdots & \phi_n(z_0) \\ \phi_0 & \phi_1(z_1) & \cdots & \phi_n(z_1) \\ \vdots & \vdots & & \vdots \\ \phi_0 & \phi_1(z_n) & \cdots & \phi_n(z_n) \end{bmatrix}$$

because  $\Phi_n(z_i) = [\phi_0, \phi_1(z_i), \dots, \phi_n(z_i)]$  and  $\phi_0 = 1$ .

- But by renormalization:

$$V_n = N_n \mathcal{V}_n, \quad N_n = \text{diag}(\|\Phi_n(z_i)\|^{-1} : i = 0, \dots, n)$$

then  $V_n^* V_n = I_{n+1}$ .

# Szegő polynomials

- What does this  $\pi$  mean in terms of orthogonal polynomials?
- Szegő OPUC:  $[1, z, z^2, z^3, \dots] \xrightarrow{\perp} [\phi_0, \phi_1, \phi_2, \dots]$   
CMV OLPU:  $[1, z, z^{-1}, z^2, z^{-2}, z^3, \dots] \xrightarrow{\perp} [\varphi_0, \varphi_1, \varphi_2, \dots]$   
 $\varphi_{2k} \in \Pi_{-k,k}, \quad \varphi_{2k+1} \in \Pi_{-k,k+1}, \quad k = 0, 1, 2, \dots$   
and  
 $\varphi_{2k}(z) = \varepsilon_{2k}[z^{-k}\phi_{2k}(z)], \quad \varphi_{2k+1}(z) = \varepsilon_{2k+1}[z^{-k}\phi_{2k+1}(z)],$   
 $\varepsilon_k \in \mathbb{T}, \quad k = 0, 1, \dots$
- Thus eigenvectors  $\tilde{\Phi}_n(z_i) = [\varphi_0, \varphi_1(z_i), \dots, \varphi_n(z_i)]$  satisfy  
 $\tilde{\Phi}_n(z_i) = \Phi_n(z_i)E_n, \quad E_n$  a diagonal of constants of modulus 1  
because  $\varepsilon_k \in \mathbb{T}$  and  $z_i^k \in \mathbb{T}$ .
- This explains why reordering the unitary  $G_k$  factors does not change the absolute values of the entries in eigenvectors.

# Szegő polynomials

- What does this  $\pi$  mean in terms of orthogonal polynomials?
- Szegő OPUC:  $[1, z, z^2, z^3, \dots] \xrightarrow{\perp} [\phi_0, \phi_1, \phi_2, \dots]$   
CMV OLPUC:  $[1, z, z^{-1}, z^2, z^{-2}, z^3, \dots] \xrightarrow{\perp} [\varphi_0, \varphi_1, \varphi_2, \dots]$   
 $\varphi_{2k} \in \Pi_{-k,k}, \quad \varphi_{2k+1} \in \Pi_{-k,k+1}, \quad k = 0, 1, 2, \dots$   
and  
 $\varphi_{2k}(z) = \varepsilon_{2k}[z^{-k}\phi_{2k}(z)], \quad \varphi_{2k+1}(z) = \varepsilon_{2k+1}[z^{-k}\phi_{2k+1}(z)],$   
 $\varepsilon_k \in \mathbb{T}, \quad k = 0, 1, \dots$
- Thus eigenvectors  $\tilde{\Phi}_n(z_i) = [\varphi_0, \varphi_1(z_i), \dots, \varphi_n(z_i)]$  satisfy  
 $\tilde{\Phi}_n(z_i) = \Phi_n(z_i)E_n, \quad E_n$  a diagonal of constants of modulus 1  
because  $\varepsilon_k \in \mathbb{T}$  and  $z_i^k \in \mathbb{T}$ .
- This explains why reordering the **unitary**  $G_k$  factors does not change the absolute values of the entries in eigenvectors.

# Szegő polynomials

- What does this  $\pi$  mean in terms of orthogonal polynomials?
- Szegő OPUC:  $[1, z, z^2, z^3, \dots] \xrightarrow{\perp} [\phi_0, \phi_1, \phi_2, \dots]$   
CMV OLPU:  $[1, z, z^{-1}, z^2, z^{-2}, z^3, \dots] \xrightarrow{\perp} [\varphi_0, \varphi_1, \varphi_2, \dots]$   
 $\varphi_{2k} \in \Pi_{-k,k}, \quad \varphi_{2k+1} \in \Pi_{-k,k+1}, \quad k = 0, 1, 2, \dots$   
and  
 $\varphi_{2k}(z) = \varepsilon_{2k}[z^{-k}\phi_{2k}(z)], \quad \varphi_{2k+1}(z) = \varepsilon_{2k+1}[z^{-k}\phi_{2k+1}(z)],$   
 $\varepsilon_k \in \mathbb{T}, \quad k = 0, 1, \dots$
- Thus eigenvectors  $\tilde{\Phi}_n(z_i) = [\varphi_0, \varphi_1(z_i), \dots, \varphi_n(z_i)]$  satisfy  
 $\tilde{\Phi}_n(z_i) = \Phi_n(z_i)E_n, \quad E_n$  a diagonal of constants of modulus 1  
because  $\varepsilon_k \in \mathbb{T}$  and  $z_i^k \in \mathbb{T}$ .
- This explains why reordering the unitary  $G_k$  factors does not change the absolute values of the entries in eigenvectors.

- What does this  $\pi$  mean in terms of orthogonal polynomials?
- Szegő OPUC:  $[1, z, z^2, z^3, \dots] \xrightarrow{\perp} [\phi_0, \phi_1, \phi_2, \dots]$   
CMV OLPU:  $[1, z, z^{-1}, z^2, z^{-2}, z^3, \dots] \xrightarrow{\perp} [\varphi_0, \varphi_1, \varphi_2, \dots]$   
 $\varphi_{2k} \in \Pi_{-k,k}, \quad \varphi_{2k+1} \in \Pi_{-k,k+1}, \quad k = 0, 1, 2, \dots$   
and  
 $\varphi_{2k}(z) = \varepsilon_{2k}[z^{-k}\phi_{2k}(z)], \quad \varphi_{2k+1}(z) = \varepsilon_{2k+1}[z^{-k}\phi_{2k+1}(z)],$   
 $\varepsilon_k \in \mathbb{T}, \quad k = 0, 1, \dots$
- Thus eigenvectors  $\tilde{\Phi}_n(z_i) = [\varphi_0, \varphi_1(z_i), \dots, \varphi_n(z_i)]$  satisfy  
 $\tilde{\Phi}_n(z_i) = \Phi_n(z_i)E_n, \quad E_n$  a diagonal of constants of modulus 1  
because  $\varepsilon_k \in \mathbb{T}$  and  $z_i^k \in \mathbb{T}$ .
- This explains why reordering the **unitary**  $G_k$  factors does not change the absolute values of the entries in eigenvectors.

- Mystery solved for  $M_\pi$  matrices.

But where do the  $A$  and  $D$  matrices come from?

- OPUC  $\rightarrow$  ORFUC (orthogonal rational functions on  $\mathbb{T}$ )

$$\alpha_0 = 0, \alpha_1, \alpha_2, \dots, \quad \alpha_k \in \mathbb{D}$$

$$B_0 = 1, B_k(z) = \prod_{i=1}^k \frac{z - \alpha_i}{1 - \bar{\alpha}_i z}, \quad k = 1, 2, 3, \dots$$

ORFUC = OPUC if all  $\alpha_k = 0$

- $[B_0, B_1, B_2, \dots] \xrightarrow{\perp} R = [\rho_0, \rho_1, \rho_2, \dots]$

- OPUC:  $\Phi(z) [z\mathcal{I} - \mathcal{G}] = 0$

$$\text{ORFUC: } R(z)S^{-1}[(z\mathcal{I} - \mathcal{A}) - (\mathcal{I} - \mathcal{A}^*z)\mathcal{G}'] = 0$$

$$\mathcal{A} = \text{diag}(\alpha_0, \alpha_1, \alpha_2, \dots), \quad S = S_{\mathcal{A}} = (\mathcal{I} - \mathcal{A}^*\mathcal{A})^{1/2}$$

$$\mathcal{G}' = G'_1 G'_2 G'_3 \cdots, \quad G'_k \text{ unitary}$$

- Mystery solved for  $M_\pi$  matrices.  
But where do the  $A$  and  $D$  matrices come from?
- OPUC  $\rightarrow$  ORFUC (orthogonal rational functions on  $\mathbb{T}$ )

$$\alpha_0 = 0, \alpha_1, \alpha_2, \dots, \quad \alpha_k \in \mathbb{D}$$

$$B_0 = 1, B_k(z) = \prod_{i=1}^k \frac{z - \alpha_i}{1 - \bar{\alpha}_i z}, \quad k = 1, 2, 3, \dots$$

$$\text{ORFUC} = \text{OPUC} \text{ if all } \alpha_k = 0$$

- $[B_0, B_1, B_2, \dots] \xrightarrow{\perp} R = [\rho_0, \rho_1, \rho_2, \dots]$
- OPUC:  $\Phi(z) [z\mathcal{I} - \mathcal{G}] = 0$

$$\text{ORFUC: } R(z)S^{-1}[(z\mathcal{I} - \mathcal{A}) - (\mathcal{I} - \mathcal{A}^*z)\mathcal{G}'] = 0$$

$$\mathcal{A} = \text{diag}(\alpha_0, \alpha_1, \alpha_2, \dots), \quad S = S_{\mathcal{A}} = (\mathcal{I} - \mathcal{A}^*\mathcal{A})^{1/2}$$

$$\mathcal{G}' = G'_1 G'_2 G'_3 \cdots, \quad G'_k \text{ unitary}$$

- Mystery solved for  $M_\pi$  matrices.  
But where do the  $A$  and  $D$  matrices come from?
- OPUC  $\rightarrow$  ORFUC (orthogonal rational functions on  $\mathbb{T}$ )

$$\alpha_0 = 0, \alpha_1, \alpha_2, \dots, \quad \alpha_k \in \mathbb{D}$$

$$B_0 = 1, B_k(z) = \prod_{i=1}^k \frac{z - \alpha_i}{1 - \bar{\alpha}_i z}, \quad k = 1, 2, 3, \dots$$

$$\text{ORFUC} = \text{OPUC} \text{ if all } \alpha_k = 0$$

- $[B_0, B_1, B_2, \dots] \xrightarrow{\perp} R = [\rho_0, \rho_1, \rho_2, \dots]$

- OPUC:  $\Phi(z) [z\mathcal{I} - \mathcal{G}] = 0$

$$\text{ORFUC: } R(z)S^{-1}[(z\mathcal{I} - \mathcal{A}) - (\mathcal{I} - \mathcal{A}^*z)\mathcal{G}'] = 0$$

$$\mathcal{A} = \text{diag}(\alpha_0, \alpha_1, \alpha_2, \dots), \quad S = S_{\mathcal{A}} = (\mathcal{I} - \mathcal{A}^*\mathcal{A})^{1/2}$$

$$\mathcal{G}' = G'_1 G'_2 G'_3 \cdots, \quad G'_k \text{ unitary}$$



- Mystery solved for  $M_\pi$  matrices.  
But where do the  $A$  and  $D$  matrices come from?
- OPUC  $\rightarrow$  ORFUC (orthogonal rational functions on  $\mathbb{T}$ )

$$\alpha_0 = 0, \alpha_1, \alpha_2, \dots, \quad \alpha_k \in \mathbb{D}$$

$$B_0 = 1, B_k(z) = \prod_{i=1}^k \frac{z - \alpha_i}{1 - \bar{\alpha}_i z}, \quad k = 1, 2, 3, \dots$$

$$\text{ORFUC} = \text{OPUC} \text{ if all } \alpha_k = 0$$

- $[B_0, B_1, B_2, \dots] \xrightarrow{\perp} R = [\rho_0, \rho_1, \rho_2, \dots]$
- OPUC:  $\Phi(z) [z\mathcal{I} - \mathcal{G}] = 0$

$$\text{ORFUC: } R(z)S^{-1}[(z\mathcal{I} - \mathcal{A}) - (\mathcal{I} - \mathcal{A}^*z)\mathcal{G}'] = 0$$

$$\mathcal{A} = \text{diag}(\alpha_0, \alpha_1, \alpha_2, \dots), \quad \mathcal{S} = \mathcal{S}_{\mathcal{A}} = (\mathcal{I} - \mathcal{A}^*\mathcal{A})^{1/2}$$

$$\mathcal{G}' = G'_1 G'_2 G'_3 \cdots, \quad G'_k \text{ unitary}$$

- $R(z)\mathcal{S}^{-1}[(z\mathcal{I} - \mathcal{A}) - (\mathcal{I} - \mathcal{A}^*z)\mathcal{G}']\mathcal{S} = 0$
- Or:  $R(z)[(z\mathcal{I} - \mathcal{A}) - (\mathcal{I} - \mathcal{A}^*z)\mathcal{G}] = 0$   
 $\mathcal{G} = \mathcal{S}^{-1}\mathcal{G}'\mathcal{S} = G_0G_1G_2\cdots$ , with  $G_k$  unitary since

$$G_k = \mathcal{S}^{-1}G'_k\mathcal{S}.$$

- $R(z)[z(\mathcal{I} + \mathcal{A}^*\mathcal{G}) - (\mathcal{G} + \mathcal{A})] = 0$
- $zR(z) = R(z) \underbrace{(\mathcal{G} + \mathcal{A})(\mathcal{I} + \mathcal{A}^*\mathcal{G})^{-1}}_{\zeta(\mathcal{G})}$
- $\mathcal{G}$  unitary  $\Rightarrow \zeta(\mathcal{G})$  (matrix Moebius transform)

- $R(z)\mathcal{S}^{-1}[(z\mathcal{I} - \mathcal{A}) - (\mathcal{I} - \mathcal{A}^*z)\mathcal{G}']\mathcal{S} = 0$
- Or:  $R(z)[(z\mathcal{I} - \mathcal{A}) - (\mathcal{I} - \mathcal{A}^*z)\mathcal{G}] = 0$   
 $\mathcal{G} = \mathcal{S}^{-1}\mathcal{G}'\mathcal{S} = G_0G_1G_2\cdots$ , with  $G_k$  unitary since

$$G_k = \mathcal{S}^{-1}G'_k\mathcal{S}.$$

- $R(z)[z(\mathcal{I} + \mathcal{A}^*\mathcal{G}) - (\mathcal{G} + \mathcal{A})] = 0$
- $zR(z) = R(z) \underbrace{(\mathcal{G} + \mathcal{A})(\mathcal{I} + \mathcal{A}^*\mathcal{G})^{-1}}_{\zeta(\mathcal{G})}$
- $\mathcal{G}$  unitary  $\Rightarrow \zeta(\mathcal{G})$  (matrix Moebius transform)

- $R(z)\mathcal{S}^{-1}[(z\mathcal{I} - \mathcal{A}) - (\mathcal{I} - \mathcal{A}^*z)\mathcal{G}']\mathcal{S} = 0$
- Or:  $R(z)[(z\mathcal{I} - \mathcal{A}) - (\mathcal{I} - \mathcal{A}^*z)\mathcal{G}] = 0$   
 $\mathcal{G} = \mathcal{S}^{-1}\mathcal{G}'\mathcal{S} = G_0G_1G_2\cdots$ , with  $G_k$  unitary since

$$G_k = \mathcal{S}^{-1}G'_k\mathcal{S}.$$

- $R(z)[z(\mathcal{I} + \mathcal{A}^*\mathcal{G}) - (\mathcal{G} + \mathcal{A})] = 0$
- $zR(z) = R(z) \underbrace{(\mathcal{G} + \mathcal{A})(\mathcal{I} + \mathcal{A}^*\mathcal{G})^{-1}}_{\zeta(\mathcal{G})}$
- $\mathcal{G}$  unitary  $\Rightarrow \zeta(\mathcal{G})$  (matrix Moebius transform)

- $R(z)\mathcal{S}^{-1}[(z\mathcal{I} - \mathcal{A}) - (\mathcal{I} - \mathcal{A}^*z)\mathcal{G}']\mathcal{S} = 0$
- Or:  $R(z)[(z\mathcal{I} - \mathcal{A}) - (\mathcal{I} - \mathcal{A}^*z)\mathcal{G}] = 0$   
 $\mathcal{G} = \mathcal{S}^{-1}\mathcal{G}'\mathcal{S} = G_0G_1G_2\cdots$ , with  $G_k$  unitary since

$$G_k = \mathcal{S}^{-1}G'_k\mathcal{S}.$$

- $R(z)[z(\mathcal{I} + \mathcal{A}^*\mathcal{G}) - (\mathcal{G} + \mathcal{A})] = 0$
- $zR(z) = R(z) \underbrace{(\mathcal{G} + \mathcal{A})(\mathcal{I} + \mathcal{A}^*\mathcal{G})^{-1}}_{\zeta(\mathcal{G})}$
- $\mathcal{G}$  unitary  $\Rightarrow \zeta(\mathcal{G})$  (matrix Moebius transform)

- $R(z)\mathcal{S}^{-1}[(z\mathcal{I} - \mathcal{A}) - (\mathcal{I} - \mathcal{A}^*z)\mathcal{G}']\mathcal{S} = 0$
- Or:  $R(z)[(z\mathcal{I} - \mathcal{A}) - (\mathcal{I} - \mathcal{A}^*z)\mathcal{G}] = 0$   
 $\mathcal{G} = \mathcal{S}^{-1}\mathcal{G}'\mathcal{S} = G_0G_1G_2\cdots$ , with  $G_k$  unitary since

$$G_k = \mathcal{S}^{-1}G'_k\mathcal{S}.$$

- $R(z)[z(\mathcal{I} + \mathcal{A}^*\mathcal{G}) - (\mathcal{G} + \mathcal{A})] = 0$
- $zR(z) = R(z) \underbrace{(\mathcal{G} + \mathcal{A})(\mathcal{I} + \mathcal{A}^*\mathcal{G})^{-1}}_{\zeta(\mathcal{G})}$
- $\mathcal{G}$  unitary  $\Rightarrow \zeta(\mathcal{G})$  (matrix Moebius transform)

- $zR(z) = R(z) \underbrace{(\mathcal{G} + \mathcal{A})(\mathcal{I} + \mathcal{A}^*\mathcal{G})^{-1}}_{\zeta(\mathcal{G})}$

- Truncate:

$$zR_n(z) = R_n(z) \underbrace{(\mathcal{G}_n + A_n)(I_{n+1} + A_n^*\mathcal{G}_n)^{-1}}_{\zeta_n(\mathcal{G}_n)} + [0, \dots, 0, c\rho_{n+1}(z)]$$

$$P = P_{n+1}^\infty, R_n = RP^*, A_n = P\mathcal{A}P^*, \mathcal{G}_n = P\mathcal{G}P^*$$

- $z = z_i$  is zero of  $\rho_{n+1}$  then it is eigenvalue of  $\zeta_n(\mathcal{G}_n)$   
and  $R_n(z_i) = [\rho_0, \rho_1(z_i), \rho_2(z_i), \dots]$  is a corresponding eigenvector.
- For a unitary truncation  $\mathcal{G}_n$  and  $\zeta_n(\mathcal{G}_n)$  are unitary

- $$zR(z) = R(z) \underbrace{(\mathcal{G} + \mathcal{A})(\mathcal{I} + \mathcal{A}^*\mathcal{G})^{-1}}_{\zeta(\mathcal{G})}$$

- Truncate:

$$zR_n(z) = R_n(z) \underbrace{(\mathcal{G}_n + A_n)(I_{n+1} + A_n^*\mathcal{G}_n)^{-1}}_{\zeta_n(\mathcal{G}_n)} + [0, \dots, 0, c\rho_{n+1}(z)]$$

$$P = P_{n+1}^\infty, R_n = RP^*, A_n = P\mathcal{A}P^*, \mathcal{G}_n = P\mathcal{G}P^*$$

- $z = z_i$  is zero of  $\rho_{n+1}$  then it is eigenvalue of  $\zeta_n(\mathcal{G}_n)$   
and  $R_n(z_i) = [\rho_0, \rho_1(z_i), \rho_2(z_i), \dots]$  is a corresponding eigenvector.
- For a unitary truncation  $\mathcal{G}_n$  and  $\zeta_n(\mathcal{G}_n)$  are unitary



- $$zR(z) = R(z) \underbrace{(\mathcal{G} + \mathcal{A})(\mathcal{I} + \mathcal{A}^*\mathcal{G})^{-1}}_{\zeta(\mathcal{G})}$$

- Truncate:

$$zR_n(z) = R_n(z) \underbrace{(\mathcal{G}_n + A_n)(I_{n+1} + A_n^*\mathcal{G}_n)^{-1}}_{\zeta_n(\mathcal{G}_n)} + [0, \dots, 0, c\rho_{n+1}(z)]$$

$$P = P_{n+1}^\infty, \quad R_n = RP^*, \quad A_n = P\mathcal{A}P^*, \quad \mathcal{G}_n = P\mathcal{G}P^*$$

- $z = z_i$  is zero of  $\rho_{n+1}$  then it is eigenvalue of  $\zeta_n(\mathcal{G}_n)$   
and  $R_n(z_i) = [\rho_0, \rho_1(z_i), \rho_2(z_i), \dots]$  is a corresponding eigenvector.
- For a unitary truncation  $\mathcal{G}_n$  and  $\zeta_n(\mathcal{G}_n)$  are unitary

- $$zR(z) = R(z) \underbrace{(\mathcal{G} + \mathcal{A})(\mathcal{I} + \mathcal{A}^*\mathcal{G})^{-1}}_{\zeta(\mathcal{G})}$$

- Truncate:

$$zR_n(z) = R_n(z) \underbrace{(\mathcal{G}_n + A_n)(I_{n+1} + A_n^*\mathcal{G}_n)^{-1}}_{\zeta_n(\mathcal{G}_n)} + [0, \dots, 0, c\rho_{n+1}(z)]$$

$$P = P_{n+1}^\infty, R_n = RP^*, A_n = P\mathcal{A}P^*, \mathcal{G}_n = P\mathcal{G}P^*$$

- $z = z_i$  is zero of  $\rho_{n+1}$  then it is eigenvalue of  $\zeta_n(\mathcal{G}_n)$   
and  $R_n(z_i) = [\rho_0, \rho_1(z_i), \rho_2(z_i), \dots]$  is a corresponding eigenvector.
- For a **unitary** truncation  $\mathcal{G}_n$  and  $\zeta_n(\mathcal{G}_n)$  are unitary

- Can we reorder the factors  $G_0 G_1 G_2 \cdots = \mathcal{G}_n$  and keep  $\sigma$  in  $\zeta_n(\mathcal{G}_n) := (\mathcal{G}_n + A_n)(I_{n+1} + A_n^* \mathcal{G}_n)^{-1}$  ?

- Can AMPD be generalized to Rational AMPD (RAMPD)?

- RAMPD:  $(AM_\pi + C)(BM_\pi + D)^{-1}$ ,  $A, B, C, D$  diagonal

- More general: pencils  $(AM_\pi + C, BM_\pi + D)$

generalized eigenvalue problem:

characteristic polynomial = determinant of  $(AM_\pi + C) - (BM_\pi + D)\lambda$

or  $\underbrace{(A - B\lambda)}_{A'} M_\pi + \underbrace{(C - D\lambda)}_{D'} \equiv A' M_\pi + D'$ ,

$A' M_\pi + D'$  is AMPD  $\Rightarrow$  determinant independent of permutation  $\pi$   
zeros of determinant are eigenvalues  $\lambda$ .

- Can we reorder the factors  $G_0 G_1 G_2 \cdots = \mathcal{G}_n$  and keep  $\sigma$  in  $\zeta_n(\mathcal{G}_n) := (\mathcal{G}_n + A_n)(I_{n+1} + A_n^* \mathcal{G}_n)^{-1}$  ?
- Can AMPD be generalized to Rational AMPD (RAMPD)?
- RAMPD:  $(AM_\pi + C)(BM_\pi + D)^{-1}$ ,  $A, B, C, D$  diagonal
- More general: pencils  $(AM_\pi + C, BM_\pi + D)$   
 generalized eigenvalue problem:  
 characteristic polynomial = determinant of  $(AM_\pi + C) - (BM_\pi + D)\lambda$   
 or  $\underbrace{(A - B\lambda)}_{A'} M_\pi + \underbrace{(C - D\lambda)}_{D'} \equiv A' M_\pi + D'$ ,  
 $A' M_\pi + D'$  is AMPD  $\Rightarrow$  determinant independent of permutation  $\pi$   
 zeros of determinant are eigenvalues  $\lambda$ .

- Can we reorder the factors  $G_0 G_1 G_2 \cdots = \mathcal{G}_n$  and keep  $\sigma$  in  $\zeta_n(\mathcal{G}_n) := (\mathcal{G}_n + A_n)(I_{n+1} + A_n^* \mathcal{G}_n)^{-1}$  ?
- Can AMPD be generalized to Rational AMPD (RAMPD)?
- RAMPD:  $(AM_\pi + C)(BM_\pi + D)^{-1}$ ,  $A, B, C, D$  diagonal
- More general: pencils  $(AM_\pi + C, BM_\pi + D)$   
 generalized eigenvalue problem:  
 characteristic polynomial = determinant of  $(AM_\pi + C) - (BM_\pi + D)\lambda$   
 or  $\underbrace{(A - B\lambda)}_{A'} M_\pi + \underbrace{(C - D\lambda)}_{D'} \equiv A' M_\pi + D'$ ,  
 $A' M_\pi + D'$  is AMPD  $\Rightarrow$  determinant independent of permutation  $\pi$   
 zeros of determinant are eigenvalues  $\lambda$ .

- Can we reorder the factors  $G_0 G_1 G_2 \cdots = \mathcal{G}_n$  and keep  $\sigma$  in  $\zeta_n(\mathcal{G}_n) := (\mathcal{G}_n + A_n)(I_{n+1} + A_n^* \mathcal{G}_n)^{-1}$  ?
- Can AMPD be generalized to Rational AMPD (RAMPD)?
- RAMPD:  $(AM_\pi + C)(BM_\pi + D)^{-1}$ ,  $A, B, C, D$  diagonal
- More general: pencils  $(AM_\pi + C, BM_\pi + D)$   
 generalized eigenvalue problem:  
 characteristic polynomial = determinant of  $(AM_\pi + C) - (BM_\pi + D)\lambda$   
 or  $\underbrace{(A - B\lambda)}_{A'} M_\pi + \underbrace{(C - D\lambda)}_{D'} \equiv A' M_\pi + D'$ ,  
 $A' M_\pi + D'$  is AMPD  $\Rightarrow$  determinant independent of permutation  $\pi$   
 zeros of determinant are eigenvalues  $\lambda$ .

- Can AMPD be generalized to Rational AMPD (RAMPD)?
- Yes we can!

## Theorem

The spectrum of the RAMPD does not depend on  $\pi$ :  
 $\sigma(AM_\pi + C, BM_\pi + D)$  independent of the permutation  $\pi$ .

- Can AMPD be generalized to Rational AMPD (RAMPD)?
- Yes we can!

## Theorem

The spectrum of the RAMPD does not depend on  $\pi$ :  
 $\sigma(AM_\pi + C, BM_\pi + D)$  independent of the permutation  $\pi$ .



- Can AMPD be generalized to Rational AMPD (RAMPD)?
- Yes we can!

## Theorem

The spectrum of the RAMPD does not depend on  $\pi$ :  
 $\sigma(AM_\pi + C, BM_\pi + D)$  independent of the permutation  $\pi$ .

- What does a different  $\pi$  means for ORF?

- Previously: if

$$[B_0, B_1, B_2, \dots] \xrightarrow{\perp} R = [\rho_0, \rho_1, \rho_2, \dots] \text{ then } R(z)[z\mathcal{I} - \zeta(\mathcal{G})] = 0 \text{ with } \mathcal{G} = G_0 G_1 G_2 \dots$$

- Now: if  $[B_0, B_1^{-1}, B_2, B_2^{-1}, \dots] \xrightarrow{\perp} \tilde{R} = [\varrho_0, \varrho_1, \varrho_2, \dots]$  then

$$\varrho_{2k} = \varepsilon_{2k} [B_k^{-1} \rho_k], \quad \varrho_{2k+1} = \varepsilon_{2k+1} [B_k^{-1} \rho_{2k+1}]$$

and

$$\tilde{R}(z)[z\mathcal{I} - \zeta(\tilde{\mathcal{G}})] = 0 \text{ with}$$

$$\tilde{\mathcal{G}} = G_0 (G_2 G_1) (G_4 G_3) \dots = (G_0 G_2 \dots) (G_1 G_3 \dots)$$

- Recall

$$B_k = \prod_{i=1}^k \frac{z - \alpha_i}{1 - \bar{\alpha}_i z} \quad \text{introduce successive poles } \frac{1}{\bar{\alpha}_i} \notin \mathbb{D}$$

$$B_k^{-1} = \prod_{i=1}^k \frac{1 - \bar{\alpha}_i z}{z - \alpha_i} \quad \text{introduce successive poles } \alpha_i \in \mathbb{D}$$

- What does a different  $\pi$  means for ORF?

- Previously: if

$$[B_0, B_1, B_2, \dots] \xrightarrow{\perp} R = [\rho_0, \rho_1, \rho_2, \dots] \text{ then } R(z)[z\mathcal{I} - \zeta(\mathcal{G})] = 0 \text{ with } \mathcal{G} = G_0 G_1 G_2 \dots$$

- Now: if  $[B_0, B_1^{-1}, B_2, B_2^{-1}, \dots] \xrightarrow{\perp} \tilde{R} = [\varrho_0, \varrho_1, \varrho_2, \dots]$  then

$$\varrho_{2k} = \varepsilon_{2k} [B_k^{-1} \rho_k], \quad \varrho_{2k+1} = \varepsilon_{2k+1} [B_k^{-1} \rho_{2k+1}]$$

and

$$\tilde{R}(z)[z\mathcal{I} - \zeta(\tilde{\mathcal{G}})] = 0 \text{ with}$$

$$\tilde{\mathcal{G}} = G_0 (G_2 G_1) (G_4 G_3) \dots = (G_0 G_2 \dots) (G_1 G_3 \dots)$$

- Recall

$$B_k = \prod_{i=1}^k \frac{z - \alpha_i}{1 - \bar{\alpha}_i z} \quad \text{introduce successive poles } \frac{1}{\bar{\alpha}_i} \notin \mathbb{D}$$

$$B_k^{-1} = \prod_{i=1}^k \frac{1 - \bar{\alpha}_i z}{z - \alpha_i} \quad \text{introduce successive poles } \alpha_i \in \mathbb{D}$$

- What does a different  $\pi$  means for ORF?

- Previously: if

$$[B_0, B_1, B_2, \dots] \xrightarrow{\perp} R = [\rho_0, \rho_1, \rho_2, \dots] \text{ then } R(z)[z\mathcal{I} - \zeta(\mathcal{G})] = 0 \text{ with } \mathcal{G} = G_0 G_1 G_2 \dots$$

- Now: if  $[B_0, B_1^{-1}, B_2, B_2^{-1}, \dots] \xrightarrow{\perp} \tilde{R} = [\varrho_0, \varrho_1, \varrho_2, \dots]$  then

$$\varrho_{2k} = \varepsilon_{2k} [B_k^{-1} \rho_k], \quad \varrho_{2k+1} = \varepsilon_{2k+1} [B_k^{-1} \rho_{2k+1}]$$

and

$$\tilde{R}(z)[z\mathcal{I} - \zeta(\tilde{\mathcal{G}})] = 0 \text{ with}$$

$$\tilde{\mathcal{G}} = G_0(G_2 G_1)(G_4 G_3) \dots = (G_0 G_2 \dots)(G_1 G_3 \dots)$$

- Recall

$$B_k = \prod_{i=1}^k \frac{z - \alpha_i}{1 - \bar{\alpha}_i z} \quad \text{introduce successive poles } \frac{1}{\bar{\alpha}_i} \notin \mathbb{D}$$

$$B_k^{-1} = \prod_{i=1}^k \frac{1 - \bar{\alpha}_i z}{z - \alpha_i} \quad \text{introduce successive poles } \alpha_i \in \mathbb{D}$$

- What does a different  $\pi$  means for ORF?
- Previously: if  $[B_0, B_1, B_2, \dots] \xrightarrow{\perp} R = [\rho_0, \rho_1, \rho_2, \dots]$  then  $R(z)[z\mathcal{I} - \zeta(\mathcal{G})] = 0$  with  $\mathcal{G} = G_0 G_1 G_2 \dots$
- Now: if  $[B_0, B_1^{-1}, B_2, B_2^{-1}, \dots] \xrightarrow{\perp} \tilde{R} = [\varrho_0, \varrho_1, \varrho_2, \dots]$  then  $\varrho_{2k} = \varepsilon_{2k}[B_k^{-1} \rho_k]$ ,  $\varrho_{2k+1} = \varepsilon_{2k+1}[B_k^{-1} \rho_{2k+1}]$  and  $\tilde{R}(z)[z\mathcal{I} - \zeta(\tilde{\mathcal{G}})] = 0$  with  $\tilde{\mathcal{G}} = G_0(G_2 G_1)(G_4 G_3) \dots = (G_0 G_2 \dots)(G_1 G_3 \dots)$
- Recall

$$B_k = \prod_{i=1}^k \frac{z - \alpha_i}{1 - \bar{\alpha}_i z} \quad \text{introduce successive poles } \frac{1}{\bar{\alpha}_i} \notin \mathbb{D}$$

$$B_k^{-1} = \prod_{i=1}^k \frac{1 - \bar{\alpha}_i z}{z - \alpha_i} \quad \text{introduce successive poles } \alpha_i \in \mathbb{D}$$

- It was first shown in the unitary case for alternating (CMV)  
 $\pi = (0, 1, 3, 2, 5, 4, \dots)$  case that  
 $\sigma(\zeta(\tilde{\mathcal{G}})) = \sigma(\zeta(\mathcal{G}))$  and after unitary truncation  
 $\sigma(\zeta_n(\tilde{\mathcal{G}}_n)) = \sigma(\zeta_n(\mathcal{G}_n))$   
 by Velázquez (2008)
- [A.B., R. Cruz-Barroso, A. Lasarow (2017)]  
 shows that this holds also for the non-unitary RAMPD for any  $\pi$ .  
 In unitary case moreover the absolute values of eigenvectors  $|R(z_i)|$   
 independent of  $\pi$ .

- It was first shown in the unitary case for alternating (CMV)  $\pi = (0, 1, 3, 2, 5, 4, \dots)$  case that  $\sigma(\zeta(\tilde{\mathcal{G}})) = \sigma(\zeta(\mathcal{G}))$  and after unitary truncation  $\sigma(\zeta_n(\tilde{\mathcal{G}}_n)) = \sigma(\zeta_n(\mathcal{G}_n))$  by Velázquez (2008)
- [A.B., R. Cruz-Barroso, A. Lasarow (2017)] shows that this holds also for the non-unitary RAMPD for any  $\pi$ . In unitary case moreover the absolute values of eigenvectors  $|R(z_i)|$  independent of  $\pi$ .



M.J. Cantero, R. Cruz-Barroso, P. González-Vera

A matrix approach to the computation of quadrature formulas on the unit circle

*Appl. Numerical Math.* 58:296–318, 2008.



L. Velázquez.

Spectral methods for orthogonal rational functions

*J. Functional Anal.*, 254(2):954–986, 2008.



A. Bultheel, R. Cruz-Barroso, A. Lasarow.

Orthogonal rational functions on the unit circle with prescribed poles not on the unit circle

*SIGMA.*, 13:1-49, 2017.